

RECEIVED: July 31, 2014

REVISED: October 31, 2014

ACCEPTED: November 13, 2014

PUBLISHED: November 24, 2014

## Attractive holographic $c$ -functions

---

Arpan Bhattacharyya,<sup>a</sup> S. Shajidul Haque,<sup>b</sup> Vishnu Jejjala,<sup>b</sup> Suresh Nampuri<sup>b</sup>  
and Álvaro Véliz-Osorio<sup>b</sup>

<sup>a</sup>Centre for High Energy Physics, Indian Institute of Science,  
C.V. Raman Avenue, Bangalore 560012, India

<sup>b</sup>National Institute for Theoretical Physics (NITheP),  
School of Physics and Center for Theoretical Physics,  
University of Witwatersrand, WITS 2050, Johannesburg, South Africa

E-mail: [arpan@cts.iisc.ernet.in](mailto:arpan@cts.iisc.ernet.in), [Shajid.Haque@wits.ac.za](mailto:Shajid.Haque@wits.ac.za),  
[vishnu@neo.phys.wits.ac.za](mailto:vishnu@neo.phys.wits.ac.za), [nampuri@gmail.com](mailto:nampuri@gmail.com),  
[Alvaro.VelizOsorio@wits.ac.za](mailto:Alvaro.VelizOsorio@wits.ac.za)

ABSTRACT: Using the attractor mechanism for extremal solutions in  $\mathcal{N} = 2$  gauged supergravity, we construct a  $c$ -function that interpolates between the central charges of theories at ultraviolet and infrared conformal fixed points corresponding to anti-de Sitter geometries. The  $c$ -function we obtain is couched purely in terms of bulk quantities and connects two different dimensional CFTs at the stable conformal fixed points under the RG flow.

KEYWORDS: Gauge-gravity correspondence, AdS-CFT Correspondence, Conformal Field Models in String Theory

ARXIV EPRINT: [1407.0469](https://arxiv.org/abs/1407.0469)

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Setting up the evolution</b>	<b>3</b>
<b>3</b>	<b>The nature of the solution</b>	<b>4</b>
<b>4</b>	<b>The construction</b>	<b>7</b>
4.1	AdS <sub>3</sub> near-horizon geometry	7
4.2	AdS <sub>2</sub> near-horizon geometry	8
4.3	Comments on the $c$ -function	9
<b>5</b>	<b>Determining the constants</b>	<b>9</b>
<b>6</b>	<b>Discussion</b>	<b>10</b>

---

## 1 Introduction

When we integrate out ultraviolet degrees of freedom to obtain a low-energy effective theory, we inevitably lose information about short distance physics. The degrees of freedom become coarse-grained in the infrared. We would like to understand how this works in detail in a gravitational setting. Extremal black holes in anti-de Sitter (AdS) space supply a useful laboratory for investigating what happens to the gravitational degrees of freedom as we perform the integration. Near to the horizon of an extremal black hole in AdS<sub>5</sub>, for example, the geometry is AdS<sub>2</sub> × S<sup>3</sup>. Generically, the dimensions of the AdS spaces at the asymptopia and the near-horizon region are different. The duality between string theory on AdS geometries and the conformal field theory (CFT) resident at the boundary of the spacetime [1–3] allows us to examine the gravitational degrees of freedom carefully because the entropy of the black hole is accounted for both by the enumeration of states in the CFT corresponding to the asymptotic region and in the CFT corresponding to the near-horizon region. We should as well see that the overall number of degrees of freedom decreases as we migrate to the interior of the spacetime. The  $c$ -theorem enables us to quantify the difference between the central charges of the two fixed point theories at either end of the flow. In this paper, we wish to make this statement precise in non-trivial supergravity environments with matter fields for which the dimension of the AdS factor changes as we traverse the radial direction in the bulk toward an extremal horizon. We will do this by providing an algorithm for constructing a  $c$ -function.

In investigating critical models in two dimensions, Zamolodchikov proved a remarkable result: when conformal fixed points are connected by a renormalization group (RG) flow, there exists a positive real function  $c(g_i, \Lambda)$  of the coupling constants and energy scale

whose value does not increase along the trajectory. At the fixed points of the flow, where the beta functions vanish, this  $c$ -function is stationary and assumes values equal to the central charges of the corresponding conformal field theories (CFT) [4]. As  $c_{UV} \geq c_{IR}$ , the RG evolution of the  $c$ -function is a gradient flow in which the number of degrees of freedom decreases. In four dimensions there are two scheme independent central charges, namely  $a$  and  $c$ . In crafting a four-dimensional analogue of the  $c$ -theorem from two dimensions, we compute the vacuum expectation value of the trace of the stress-energy tensor to be

$$16\pi^2 \langle T^\mu_\mu \rangle = -a E_4 + c W_{\mu\nu\rho\sigma} W^{\mu\nu\rho\sigma}, \tag{1.1}$$

where  $E_4$ , the Euler density, is quadratic in the Riemann tensor and  $W_{\mu\nu\rho\sigma}$  is the Weyl tensor. While in all known examples,  $a_{UV} \geq a_{IR}$ , the same is not true of the coefficient  $c$  in (1.1). Thus, it is the coefficient of the Euler density that offers a candidate for the  $c$ -function in higher dimensions [5–7], and indeed, the  $a$ -theorem is now firmly established in four dimensions [8, 9]. To keep the discussion general, we will henceforth refer to the  $a$ -theorem as the  $c$ -theorem. We are referring to a function that interpolates between central charges of fixed point CFTs in a strictly monotonic manner.

The gauge/gravity duality allows us to equate the  $c$ -theorem in the boundary field theory to a statement about the holographically dual gravitational background [10–16]. The gauge/gravity duality supplies a map from the Hilbert space of states and the interactions between the various fields in the bulk gravitational background to the spectrum of states and interactions in the boundary field theory. In particular, in the context of  $AdS_5/CFT_4$ , the vacuum state of the boundary CFT is identified with empty  $AdS_5$ . An ensemble of states at finite charge density and zero temperature corresponds to an asymptotically  $AdS_5$  extremal black background while an ensemble of states at finite charge density and non-zero temperature corresponds to an asymptotically  $AdS_5$  non-extremal black background. A deformation of the boundary field theory from its ultraviolet conformal fixed point by a relevant operator initiates an RG flow that terminates at a conformal fixed point in the infrared. This RG flow produces a change in the coupling constants in the theory. The gauge/gravity duality maps this flow in the coupling constants to a radial flow of the holographically dual scalar fields in the gravitational background. The RG flow in the boundary is parameterized by a  $c$ -function, and hence the radial flow of the scalar fields will be parameterized by the holographically dual  $c$ -function in the bulk.

For spherically symmetric, static configurations in four-dimensional, two derivative gravity, a monotonically decreasing function was proposed in [17] that applies to non-vacuum settings. This is simply the area of radial slices in the bulk. At the horizon of the black hole, i.e., in the infrared, the entropy is an  $SO(3)$  invariant function that counts the degrees of freedom. This proposal is, however, unsatisfactory because it diverges at the boundary.

In this paper, we establish that for  $AdS$  spaces with single centered black holes, the  $c$ -theorem arises as a consequence of the attractor mechanism [18–26]. At large radius, the geometry is asymptotically  $AdS_{d+1}$ . Near the horizon of the black hole (black brane), the metric has an  $AdS_3$  or  $AdS_2$  factor. The  $c$ -function we construct therefore interpolates

between AdS spacetimes of *different* dimensions. At the endpoints of the flow, the  $c$ -function is extremized and assumes the values of the central charges corresponding to the AdS geometry in the ultraviolet or the infrared. Since we work in  $\mathcal{N} = 2$  supergravity, the stress-energy tensor in the bulk is highly non-trivial. Typically scalar fields and fluxes are present.

The attractor mechanism fixes the values of the scalar fields in the near-horizon geometry in terms of the quantum numbers of the black hole. The attractor values are independent of the asymptotic moduli at infinity. This is a consequence of the “deep throat” of the near-horizon AdS factor. All normalizable perturbations in the asymptotic values of the scalar fields are damped as they traverse the deep throat in the near-horizon regime. Consequently, the values of the scalars in the near-horizon region are completely independent of their values at infinity. The black hole horizon therefore acts as an attractor fixed point in the space of scalar fields. Crucially, the attractor mechanism is independent of the number of supersymmetries supported by the black hole background. In order to illustrate our argument we specialize to the attractor mechanism in a background that preserves half the supersymmetries of the theory, but this is for clarity and convenience only. Our argument readily generalizes to non-supersymmetric settings. The key to the attractor mechanism is *extremality*, not supersymmetry.

The extremal black backgrounds we consider are solutions to four-dimensional and five-dimensional  $\mathcal{N} = 2$  gauged supergravity actions that arise as consistent low-energy tree level effective actions of flux compactification in type IIB string theory. String theory in this background is holographically dual to a boundary field theory in the large- $N$  limit where the four-dimensional central charges  $a$  and  $c$  are equal. Our analysis provides a detailed recipe for computing holographic  $c$ -functions that interpolate between zero temperature states at finite charge density at the ultraviolet and infrared fixed points of the boundary field theory. These are non-vacuum states.

The outline of the paper is as follows. In section 2, we detail the attractor flows that we investigate in this work. In section 3, we develop the form of the  $c$ -function in supergravity. The null energy condition ensures that the function is monotonic. In section 4, we write the  $c$ -functions explicitly for black backgrounds with  $\text{AdS}_{d+1}$  asymptopia and an  $\text{AdS}_3$  or an  $\text{AdS}_2$  factor in the near-horizon region. In section 5, we evaluate the example of a black brane in  $\text{AdS}_5$  with a near-horizon  $\text{AdS}_3$  factor and explain how to fix the constants in the  $c$ -function. In section 6, we discuss our results and provide a prospectus for future work.

## 2 Setting up the evolution

In holographic RG, the radial direction in AdS is a proxy for the scale of the Wilsonian flow in the dual field theory [27–30]. In the bulk, we then have a function  $c(r) = F(\phi^i(r))$ , where  $\phi^i(r)$  are scalar fields in the spacetime that are dual to the operators sourced by the coupling constants on the boundary. In the dual field theory, the  $c$ -function curve is given by  $c(g_i, \Lambda) = F(g_i(\Lambda))$ , where  $\Lambda$  is the energy scale. The evolution of the  $c$ -function in the field theory follows a Callan-Symanzik equation where the choice of how we parameterize energy corresponds to the choice of the bulk radial coordinate. In describing

the  $c$ -function, we are perfectly free to think of it as a function of  $r^2$ , for instance. There is an inbuilt redundancy in the characterization of the  $c$ -function corresponding to this freedom. We need to demand only that at the endpoints of the RG flow, the function takes values corresponding to the central charges at the fixed points, where the theory is exactly conformal. These central charges are related to the radius of the AdS geometry dual to the fixed point. In between the endpoints, due to the choice of the radial coordinate, the flow can be different, but each of the  $c$ -functions obtained in this manner is monotonic.

We specialize to  $\mathcal{N} = 2$  U(1) gauged supergravity and use the attractor mechanism for black solutions in the bulk [22–26]. The Hilbert space of the dual field theory is graded by temperature, charge, angular momentum, and flux quantum numbers that also distinguish the black backgrounds in the bulk. As we consider single centered solutions here, we restrict the field theory to the subspace of the Hilbert space with the appropriate charges and consider the Wilsonian evolution of only this subspace. In particular, we will not consider solutions with multipole moments or hair. The states we enumerate in this subspace contribute to the universal large charge limit of the entropy. In the bulk, these states are recognized as single centered black hole (black brane) microstates, whose number reproduces the Bekenstein-Hawking entropy. For extremal black holes, the entropy function  $S = \frac{1}{4}A$  (in units where the Newton constant  $G = 1$  and  $A$  is the area of the horizon) is a fixed quantity.

The near-horizon geometry of the extremal background is of the form  $\text{AdS}_2 \times X$  or  $\text{AdS}_3 \times X$ , where  $X$  can be spherical, hyperbolic, or planar depending on the structure of the black hole (black brane) horizon. The horizon acts as an attractive fixed point for the scalar flows in this background. An attractive on-shell scalar flow from the asymptotic AdS boundary to the near-horizon AdS factor is holographically dual to the Wilsonian flow of the effective field theory action that encodes the dynamics of the field theory operators dual to the scalar fields in the specified black background. The flow interpolates between the ultraviolet fixed point CFT corresponding to the asymptotic AdS geometry and the infrared fixed point CFT corresponding to the near-horizon AdS geometry. The near-horizon AdS geometry is dual either to a two-dimensional CFT or to the discrete light-cone quantization (DLCQ) limit of one [31]. The subset of states that corresponds to the single centered black solution are encoded in a Virasoro algebra representation of a two-dimensional CFT in the infrared. The  $c$ -function that acts as an affine parameter for this Wilsonian flow is dual to an affine parametrization of the attractive scalar flows in the bulk.  $\text{AdS}_3 \times X$  is stable under small perturbations. Infrared fixed points of the form of  $\text{AdS}_2 \times X$  have been studied extensively especially in the context of stability under perturbations and the interested reader is referred to [32–36] for further details.

### 3 The nature of the solution

In order to focus our discussion, we recall that in four-dimensional  $\mathcal{N} = 2$  gauged supergravity in the presence of fluxes that arise from a low-energy string compactification, the gravity multiplet couples to massless gauge fields. The coupling strengths of these interactions are determined by the scalars that we have mentioned. While supersymmetry will not

ultimately turn out to be essential to the analysis, we concentrate initially on supersymmetric flows in asymptotically AdS, static, spherically symmetric, extremal backgrounds. Deducing a suitable  $c$ -function in the bulk is equivalent to finding a real function of the scalar fields which is invariant under symplectic transformations of the scalar moduli space. We seek to encode this proposal in terms of purely geometrical quantities and then to generalize the result to non-supersymmetric attractive flows in any dimension. The central charges are read off at both boundaries by computing the renormalized part of the stress-energy tensor. These terms depend purely on the metric and are independent of the matter content of the theory. This motivates the idea that something purely geometrical arrests all the information about the central charges at the fixed points of the flow. The  $c$ -function must have extremization conditions at these endpoints and monotonicity properties that are independent of the matter content of the field theory and must depend on the Einstein tensor  $G_{\mu\nu}$ .

To ensure monotonicity of the  $c$ -function, we examine its derivative. A geometric function  $c(r)$  must have a derivative  $c'(r)$  that is proportional to a scalar quantity in gravity constructed purely out of the stress-energy tensor with a definite signature. (All primes denote differentiation with respect to the radial coordinate  $r$ .) A natural choice follows from the null energy condition, which states that for any null vector  $k$ ,  $T_{\mu\nu}k^\mu k^\nu \geq 0$  for all physical backgrounds. The inequality is saturated only in vacuum backgrounds. This is a physical input for the construction.

In order to deduce a suitable  $c$ -function, we first observe that the attractive scalar flow can be encoded in terms of a first order equation,

$$\phi^{i'} = f^i(g_{\mu\nu}(r), \phi^i(r)), \tag{3.1}$$

where  $\phi^i$ 's are the scalar fields in the bulk and  $g_{\mu\nu}$  represents the bulk metric. In the static and spherically symmetric backgrounds that we consider,  $\phi^i$  is a function only of the radial coordinate  $r$ . From (3.1), it follows that in the scalar moduli space, a vector valued function  $f(r)$  can be defined as a conservative function that vanishes at both endpoints of the flow and can be written as a gradient  $f^i(r) = G^{ij}\partial_j\Upsilon$ , where  $G_{ij}$  is the metric on the scalar space. Hence, the attractor equation can be rewritten as

$$\phi^{i'}(r) = G^{ij}\partial_j\Upsilon. \tag{3.2}$$

Furthermore, in such a first order dynamical system, there is a first order equation which relates the metric coefficients and their derivatives to the function  $\Upsilon$  that drives the scalar flow:

$$\Upsilon = p(g_{\mu\nu}(r), g'_{\mu\nu}(r)). \tag{3.3}$$

In order for  $\phi^{i'}(r)$  to vanish, the function  $\Upsilon$  has to be extremized only at the endpoints of the flow. Hence, a function that serves as an effective affine parameter for the attractive flow in moduli space could simply be proportional to  $\Upsilon$  up to a constant:

$$c(\phi^i) = \lambda + \kappa\Upsilon. \tag{3.4}$$

The function  $\Upsilon$  is proportional to the generalized superpotential in  $\mathcal{N} = 2$  theories and in particular becomes proportional to the central charges for supersymmetric attractive flows. This choice is motivated by the need to define a function which is specified by a minimum number of dynamical conditions and parameters of the system and applies our understanding that along supersymmetric attractive flows the central charge gets minimized. In order to define the bulk equivalent of the  $c$ -function for the RG flow in the boundary, we need to write down  $c$  as a function of the radial coordinate  $r$  dual to the boundary energy scale in terms of bulk geometrical quantities. The simplest procedure to do this would be to use the attractor equations to encode  $\Upsilon$  in terms of the geometry. From (3.3), we see that the function  $p(r)$  performs this encoding operation and by construction is monotonic along the attractor flow.

We write down a  $c$ -function that is monotonic as

$$c(r) = \lambda + \kappa [\chi(p(r)) + \mathcal{G}(r)], \tag{3.5}$$

where  $\mathcal{G}(r)$  is a differentiable monotonic function and  $\chi(p(r))$  is a differentiable function in  $p(r)$  which is extremized in scalar moduli space at the endpoints. We label the term in brackets in (3.5) as  $\mathcal{H}(r)$  for convenience. Imposing the absence of extremization along the flow, except at the two endpoints, leads us to the constraint that the derivative  $\mathcal{H}'(r)$  is of one sign only:

$$\mathcal{H}'(r) \geq 0 \quad \text{or} \quad \mathcal{H}'(r) \leq 0, \tag{3.6}$$

with the inequality saturated only at the endpoints  $r = r_h$  and  $r = \infty$ .

The only scalar quantity that depends on the matter content of the theory and has a definite signature determined purely from the geometry for a physical on-shell background with a null condition achieved on vacuum solutions is a contraction of the stress tensor  $T_{\mu\nu}$  with any null vector  $k^\mu$ :

$$8\pi T_{\mu\nu} k^\mu k^\nu = G_{\mu\nu} k^\mu k^\nu \geq 0. \tag{3.7}$$

Now, we can always write this contraction as

$$8\pi T_{\mu\nu} k^\mu k^\nu = \mathcal{F}(g_{\mu\nu}) \mathcal{B}(g_{\mu\nu}, g'_{\mu\nu}), \tag{3.8}$$

where  $\mathcal{F}$  is a positive regular function of the metric and  $\mathcal{B}$  is a positive regular function of the metric and its derivative. For a given  $\chi(p(r))$ , we calculate  $\mathcal{G}(r)$  by defining

$$\mathcal{H}'(r) = \frac{8\pi T_{\mu\nu} k^\mu k^\nu}{\mathcal{F}(g_{\mu\nu})}. \tag{3.9}$$

It is then manifestly true that  $\mathcal{H}'(r) \geq 0$  as a consequence of the null energy condition. We therefore determine that  $\mathcal{H}(r)$ , and consequently  $c(r)$ , is a monotonically decreasing function from the ultraviolet to the infrared. The null energy condition fixes the behavior of  $\mathcal{H}(r)$  completely. The constants  $\lambda$  and  $\kappa$  in (3.5) are determined by matching to the central charges at the two endpoints of the flow. This is our prescription for deriving the  $c$ -function.

For a given null vector and  $p(r)$ , we can in fact choose multiple functions  $\mathcal{F}(r)$  and  $\mathcal{B}(r)$  functions with the same product and, in consequence, various corresponding functions  $\chi$ . The freedom in choosing the  $\mathcal{F}$  function is related to the arbitrariness in picking a  $c$ -function for RG flows on the boundary depending on the renormalization scheme used. Furthermore, in going from the central charge to a geometrical representation, there is an extra degree of freedom available in terms of the  $SO(d-1, 1)$  group of coordinate diffeomorphisms which can operate on the non-radial part of the metric. We fix the diffeomorphism gauge by choosing the metric to be static and spherically symmetric. Such a metric can support black backgrounds with an  $AdS_2$  factor or an  $AdS_3$  factor in the infrared as the endpoint of the attractive flow.

#### 4 The construction

To be explicit, we start with a  $(d+1)$ -dimensional static spherically symmetric metric of the form

$$ds^2 = -a(r)^2 dt^2 + a(r)^{-2} dr^2 + b(r)^2 \sum_{i=1}^{d-2} dx_i^2 + w(r)^2 dz^2 . \tag{4.1}$$

The coordinate  $r$  is the radial direction corresponding to the RG flow, and  $t, x_i$ , and  $z$  are boundary coordinates. The harmonic function  $a(r)$  and the warp factors  $b(r)$  and  $w(r)$  that appear in (4.1) describe a black hole spacetime in the infrared and give an asymptotically  $AdS_{d+1}$  geometry for large  $r$ . The horizon of the extremal black hole localizes at  $r = r_h$ , where  $a(r_h) = 0$ . The singularity exists at  $b(r) = 0$ . At the horizon  $r = r_h$ , the area is written in terms of  $b(r_h) \sqrt{w(r_h)}$  [37]. Crucially, the metric (4.1) allows for a near-horizon geometry with an  $AdS_2$  factor when  $w(r) = b(r)$  and for an  $AdS_3$  factor when  $w(r) = a(r)$ .

Consider the null vector

$$k^\mu = (a(r)^{-1}, a(r), 0, \dots, 0) . \tag{4.2}$$

From the Einstein equation for the  $(d+1)$ -dimensional metric given in (4.1), the null energy contraction reads

$$8\pi T_{\mu\nu} k^\mu k^\nu = -a(r)^2 \left( \frac{w''(r)}{w(r)} + (d-2) \frac{b''(r)}{b(r)} \right) \geq 0 . \tag{4.3}$$

##### 4.1 $AdS_3$ near-horizon geometry

We set  $w(r) = a(r)$ . The attractor equations tell us that [37]

$$p(r) = \frac{a'(r)}{a(r)} + (d-2) \frac{b'(r)}{b(r)} . \tag{4.4}$$

Now, keeping in mind (3.8) and the arguments of section 3, we choose  $\mathcal{F}(r) = a(r)^2$  and let  $\chi(p(r)) = -p(r)$ . Then, in order to simultaneously satisfy

$$\mathcal{H}(r) = -p(r) + \mathcal{G}(r) \quad \text{and} \quad \mathcal{H}'(r) = \frac{8\pi T_{\mu\nu} k^\mu k^\nu}{\mathcal{F}(r)} , \tag{4.5}$$

we deduce

$$\mathcal{G}(r) = - \int dr \left\{ \left( \frac{a'(r)}{a(r)} \right)^2 + (d-2) \left( \frac{b'(r)}{b(r)} \right)^2 \right\} . \quad (4.6)$$

Putting the pieces together, the  $c$ -function can be formally written as

$$c_{\text{AdS}_3} = \lambda + \kappa \left[ \left( \frac{a'(r)}{a(r)} + (d-2) \frac{b'(r)}{b(r)} \right) + \int dr \left\{ \left( \frac{a'(r)}{a(r)} \right)^2 + (d-2) \left( \frac{b'(r)}{b(r)} \right)^2 \right\} \right] , \quad (4.7)$$

where we have absorbed signs into  $\kappa$ . We verify that  $c(r)$  as defined in (4.7) is monotonically non-increasing from the ultraviolet to the infrared as a consequence of (4.3).

## 4.2 AdS<sub>2</sub> near-horizon geometry

Similarly, using  $w(r) = b(r)$  in order to ensure the existence of an AdS<sub>2</sub> factor in the near-horizon geometry, we employ the attractor equation [39]

$$p(r) = (d-1) \frac{b'(r)}{b(r)} . \quad (4.8)$$

Following the same arguments as in section 4.1, we choose

$$\mathcal{F}(r) = (d-1) \frac{a(r)^2}{b(r)} , \quad \chi(p(r)) = - \frac{b(r)p(r)}{d-1} = -b'(r) . \quad (4.9)$$

The relations

$$\mathcal{H}(r) = -b'(r) + \mathcal{G}(r) \quad \text{and} \quad \mathcal{H}'(r) = \frac{8\pi T_{\mu\nu} k^\mu k^\nu}{\mathcal{F}(r)} \quad (4.10)$$

are satisfied for

$$\mathcal{G}(r) = 0 . \quad (4.11)$$

This yields the  $c$ -function

$$c_{\text{AdS}_2} = \lambda + \kappa b'(r) . \quad (4.12)$$

The extremization of the  $c$ -function follows from the fact that  $b''(r) = 0$  in all AdS spaces as these geometries saturate the null energy condition. One can also obtain the  $p(r)$  by looking at illustrative examples of interpolating solutions between AdS<sub>4</sub> and AdS<sub>2</sub> as in [38–40] as a consistency check.

To compare the expressions (3.4) and (4.12), let us define  $\phi^i = X^i/X^0$ , where  $X^I$  are the scalars in the vector multiplets of the  $\mathcal{N} = 2$  supergravity theory in four dimensions. (Note that  $i = 1, \dots, n_V$  and  $I = 0, \dots, n_V$ , with  $n_V + 1$  the number of vector multiplets.) Following [38, 39], we may calculate

$$b'(r) = (ab)^{-1} \mathcal{Z}(\phi) , \quad (4.13)$$

where

$$\mathcal{Z}(\phi) = |Z(\phi) - ib^2 W(\phi)| \quad (4.14)$$

is the generalized superpotential, a combination of the central charge  $Z$  and the superpotential  $W$ . With this structure, we can explicitly verify that  $b'(r_h)$  is regular. The function  $\Upsilon$  then is

$$\Upsilon = b'(r) = \frac{1}{ab} |Z(\phi) - ib^2 W(\phi)| . \quad (4.15)$$

### 4.3 Comments on the $c$ -function

The form of the  $c$ -function in (4.7) and (4.12) is robust. This function interpolates between Lorentzian fixed points in the geometry, is written purely in terms of geometrical quantities, *viz.*, the harmonic function and the warp factor in the metric, and satisfies the monotonicity and extremization properties in Zamolodchikov’s theorem. The unknown parameters are determined by the boundary conditions.

The construction in this paper therefore applies to extremal black solutions that interpolate between  $\text{AdS}_{d+1}$  asymptopia to  $\text{AdS}_2$  or  $\text{AdS}_3$  near-horizon regimes. Examples include (a) extremal black branes in five dimensions, (b) extremal black strings in five dimensions, and (c) BPS black branes and black holes in four dimensions [41]:

$$\begin{aligned}
 \text{a.} \quad & \text{AdS}_5 \xrightarrow{\text{RG}} \text{AdS}_2 \times \mathbb{R}^3, \\
 \text{b.} \quad & \text{AdS}_5 \xrightarrow{\text{RG}} \text{AdS}_3 \times \Sigma_k^2, \\
 \text{c.} \quad & \text{AdS}_4 \xrightarrow{\text{RG}} \text{AdS}_2 \times \Sigma_k^2.
 \end{aligned}
 \tag{4.16}$$

Here,  $\Sigma_k^2$  refers to two-dimensional flat space ( $k = 0$ ), the two-sphere ( $k = 1$ ), and hyperbolic space ( $k = -1$ ).

Case (c) requires some elaboration. For  $\text{AdS}_4$ , the corresponding CFT is three-dimensional. For odd dimensional CFTs, the vanishing trace anomaly term implies a vanishing central charge. In these theories, the free energy of the CFTs conformally mapped to a sphere is proposed to be the monotonically decreasing function that is stationary at the ultraviolet and infrared fixed points. More specifically, the conjectured  $c$ -function can be written as

$$F = (-1)^{\frac{d-1}{2}} \log |Z|, \quad d \text{ odd}, \tag{4.17}$$

where  $|Z|$  is the  $S^d$  partition function. For interpolating solutions between  $\text{AdS}_4$  and  $\text{AdS}_2$ , the free energy thus defined at the endpoint CFTs provides the right boundary value data to determine the constants of the bulk  $c$ -function [42, 43]. In  $\text{AdS}_2$ , the finite part of the partition function has been shown to be nothing but the dimension of the Hilbert space of the dual  $\text{CFT}_1$  [44]. This corresponds to the Hilbert space of microstates of the single centered black hole. In consequence, the boundary value data at the infrared fixed point is nothing but the entropy of the black hole or the entropy density of the black brane.

As monotonicity and extremization are consequences of the null energy condition, our proposal generalizes to all attractive flows in four dimensions including the non-supersymmetric, extremal ones.

## 5 Determining the constants

As a heuristic example, we present a model calculation to determine the constants of the  $c$ -function in an  $\text{AdS}_5$  to  $\text{AdS}_3$  interpolating black brane solution. Consistent with the normalization of [14], the central charge of the four-dimensional CFT at the boundary is given in terms of the  $\text{AdS}_5$  radius  $L_{\text{UV}}$  as

$$c_{\text{UV}} = \frac{\pi^2 L_{\text{UV}}^3}{\ell_5^3}, \tag{5.1}$$

where  $\ell_5$  is the five-dimensional Planck scale. As we are working in Einstein gravity, the  $a$  and  $c$  central charges are the same in the ultraviolet. The central charge of the infrared CFT is given in terms of the AdS<sub>3</sub> radius  $L_{\text{IR}}$  [45]:

$$c_{\text{IR}} = \frac{3L_{\text{IR}}}{2\ell_3} = \frac{3b^2(r_h)L_{\text{IR}}}{2\ell_5}, \quad (5.2)$$

where  $\ell_3$  is the three-dimensional Planck scale. (In our conventions,  $G_D = \ell_D^{D-2}$ .) Here, we can think of the planar part as a sphere of infinite radius. The two constants in the  $c$ -function are then

$$\lambda = \frac{\mathcal{H}_{\text{UV}} c_{\text{IR}} - \mathcal{H}_{\text{IR}} c_{\text{UV}}}{\mathcal{H}_{\text{UV}} - \mathcal{H}_{\text{IR}}}, \quad \kappa = \frac{c_{\text{UV}} - c_{\text{IR}}}{\mathcal{H}_{\text{UV}} - \mathcal{H}_{\text{IR}}}. \quad (5.3)$$

In the previous expression,  $\mathcal{H}_{\text{UV}}$  (respectively,  $\mathcal{H}_{\text{IR}}$ ) is the term in square brackets in (4.7) evaluated at  $r = \infty$  ( $r = r_h$ ).

Notice that, in so far as we can regard the central charge as encoding the number of degrees of freedom, at the infrared fixed point, which is a direct product of a two-dimensional space with AdS<sub>3</sub>, the number of degrees of freedom, and hence, the central charge, is formally infinite in cases where the two-dimensional space is non-compact. In these cases, one simply quotients out the phase space by the volume of the two-dimensional space to get a finite volume density. This leaves the phase space of AdS<sub>3</sub> diffeomorphisms which gives rise to a Virasoro  $\times$  Virasoro algebra unchanged as the two spaces combine as a direct product to give the infrared geometry. This is formally equivalent to taking the phase space to be the direct product of the space of diffeomorphisms of the AdS<sub>3</sub> factor and unit volume of the phase space of the trivial fiber, thus giving rise to a well-regulated finite  $c$ -function value at the infrared fixed point.

## 6 Discussion

The algorithm for determining the  $c$ -function is motivated by the general first order dynamics of the attractive flows which link the flow to a conservative vector valued function  $f(r)$  over the scalars. For a given diffeomorphism gauge, we can write down the geometrical quantity that encodes the function  $f(r)$ , and this allows us to deduce a  $c$ -function in that gauge. The null energy condition, with a careful choice of null vector, implies monotonicity; extremization follows from the scalar flow in the bulk.

The physics at the ultraviolet and infrared fixed points of the boundary theory must be independent of the choice of scheme for the RG evolution. Certainly, the  $c$ -function is not unique. Based on [46, 47], AdS/CFT equates diffeomorphism invariance in the bulk to scheme independence in the boundary theory [48]. We expect that other  $c$ -functions can be generated through general coordinate transformations in the spacetime.

Our results may be useful in investigating intermediate scaling vacua in attractor flows between critical points in moduli space. The scaling regions between two fixed points of the flow are vacuum solutions of gauged supergravity, except that they break Lorentz symmetry and correspond to points in the RG flow on the boundary where the Hilbert space that

contributes to the single centered black hole entropy contains Lorentz violating states. A class of such solutions, namely the Lifshitz geometries, has been studied extensively as candidates for field theory states that exhibit quantum critical phase transitions [39, 49, 50].

In [51], bulk and boundary flows are shown to agree at the infrared fixed point via computation of three-point correlation functions of chiral primary operators in the two-dimensional CFT. Since the  $c$ -function provides an affine parametrization of the flow globally, one use of our formalism is to generalize this result at first order away from the fixed point. Our prescription allows the construction of  $c$ -functions in higher spin theories, which provide a proving ground for essential features of holography. In these backgrounds, where the horizon for a black hole is a frame dependent concept [52], the  $c$ -function may help define an unambiguous notion of an infrared fixed point that characterizes the spacetime. As well, in the maps in (4.16), we study black solutions either in  $\text{CFT}_2$  or in  $\text{CFT}_4$ . The existence of  $c$ -functions of the type we have discussed are characterized by embeddings of the Virasoro algebra in quantum field theories with  $\text{SO}(d, 2)$  conformal symmetry upon taking certain limits. Also, it will be interesting to apply our procedure when there are higher derivative corrections [22, 40, 53–56]. One more important direction of investigation is to determine how to reproduce our proposed  $c$ -function from holographic entanglement entropy calculation [13]. We aim to explore these ideas in future work.

## Acknowledgments

We thank Paolo Benincasa, Gabriel Cardoso, Robert de Mello Koch, Kevin Goldstein, Rob Myers, Shubho Roy, Joan Simón, Aninda Sinha, and Jan Troost for deep discussions. VJ is supported by the South African Research Chairs Initiative of the Department of Science and Technology and National Research Foundation. He is grateful to KIAS for hospitality during the concluding stages of this work. SSH, SN, and AVO are supported by SARChI, NRF, and the University Research Council.

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] J.M. Maldacena, *The large- $N$  limit of superconformal field theories and supergravity*, *Int. J. Theor. Phys.* **38** (1999) 1113 [[hep-th/9711200](#)] [[INSPIRE](#)].
- [2] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from noncritical string theory*, *Phys. Lett.* **B 428** (1998) 105 [[hep-th/9802109](#)] [[INSPIRE](#)].
- [3] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)] [[INSPIRE](#)].
- [4] A.B. Zamolodchikov, *Irreversibility of the flux of the renormalization group in a 2D field theory*, *JETP Lett.* **43** (1986) 730 [[INSPIRE](#)].
- [5] J.L. Cardy, *Is there a  $c$  theorem in four-dimensions?*, *Phys. Lett.* **B 215** (1988) 749 [[INSPIRE](#)].

- [6] H. Osborn, *Derivation of a four-dimensional c theorem*, *Phys. Lett. B* **222** (1989) 97 [[INSPIRE](#)].
- [7] I. Jack and H. Osborn, *Analogs for the c theorem for four-dimensional renormalizable field theories*, *Nucl. Phys. B* **343** (1990) 647 [[INSPIRE](#)].
- [8] Z. Komargodski and A. Schwimmer, *On renormalization group flows in four dimensions*, *JHEP* **12** (2011) 099 [[arXiv:1107.3987](#)] [[INSPIRE](#)].
- [9] Z. Komargodski, *The constraints of conformal symmetry on RG flows*, *JHEP* **07** (2012) 069 [[arXiv:1112.4538](#)] [[INSPIRE](#)].
- [10] L. Girardello, M. Petrini, M. Porrati and A. Zaffaroni, *Novel local CFT and exact results on perturbations of  $N = 4$  super Yang-Mills from AdS dynamics*, *JHEP* **12** (1998) 022 [[hep-th/9810126](#)] [[INSPIRE](#)].
- [11] D.Z. Freedman, S.S. Gubser, K. Pilch and N.P. Warner, *Renormalization group flows from holography supersymmetry and a c theorem*, *Adv. Theor. Math. Phys.* **3** (1999) 363 [[hep-th/9904017](#)] [[INSPIRE](#)].
- [12] G. Lopes Cardoso, G. Dall'Agata and D. Lüst, *Curved BPS domain walls and RG flow in five-dimensions*, *JHEP* **03** (2002) 044 [[hep-th/0201270](#)] [[INSPIRE](#)].
- [13] R.C. Myers and A. Sinha, *Seeing a c-theorem with holography*, *Phys. Rev. D* **82** (2010) 046006 [[arXiv:1006.1263](#)] [[INSPIRE](#)].
- [14] R.C. Myers and A. Sinha, *Holographic c-theorems in arbitrary dimensions*, *JHEP* **01** (2011) 125 [[arXiv:1011.5819](#)] [[INSPIRE](#)].
- [15] J.T. Liu, W. Sabra and Z. Zhao, *Holographic c-theorems and higher derivative gravity*, *Phys. Rev. D* **85** (2012) 126004 [[arXiv:1012.3382](#)] [[INSPIRE](#)].
- [16] M.F. Paulos, *Holographic phase space: c-functions and black holes as renormalization group flows*, *JHEP* **05** (2011) 043 [[arXiv:1101.5993](#)] [[INSPIRE](#)].
- [17] K. Goldstein, R.P. Jena, G. Mandal and S.P. Trivedi, *A c-function for non-supersymmetric attractors*, *JHEP* **02** (2006) 053 [[hep-th/0512138](#)] [[INSPIRE](#)].
- [18] S. Ferrara, R. Kallosh and A. Strominger,  *$N = 2$  extremal black holes*, *Phys. Rev. D* **52** (1995) 5412 [[hep-th/9508072](#)] [[INSPIRE](#)].
- [19] A. Strominger, *Macroscopic entropy of  $N = 2$  extremal black holes*, *Phys. Lett. B* **383** (1996) 39 [[hep-th/9602111](#)] [[INSPIRE](#)].
- [20] S. Ferrara and R. Kallosh, *Supersymmetry and attractors*, *Phys. Rev. D* **54** (1996) 1514 [[hep-th/9602136](#)] [[INSPIRE](#)].
- [21] S. Ferrara and R. Kallosh, *Universality of supersymmetric attractors*, *Phys. Rev. D* **54** (1996) 1525 [[hep-th/9603090](#)] [[INSPIRE](#)].
- [22] A. Sen, *Black hole entropy function and the attractor mechanism in higher derivative gravity*, *JHEP* **09** (2005) 038 [[hep-th/0506177](#)] [[INSPIRE](#)].
- [23] K. Goldstein, N. Iizuka, R.P. Jena and S.P. Trivedi, *Non-supersymmetric attractors*, *Phys. Rev. D* **72** (2005) 124021 [[hep-th/0507096](#)] [[INSPIRE](#)].
- [24] D. Astefanesei, K. Goldstein, R.P. Jena, A. Sen and S.P. Trivedi, *Rotating attractors*, *JHEP* **10** (2006) 058 [[hep-th/0606244](#)] [[INSPIRE](#)].

- [25] A. Dabholkar, A. Sen and S.P. Trivedi, *Black hole microstates and attractor without supersymmetry*, *JHEP* **01** (2007) 096 [[hep-th/0611143](#)] [[INSPIRE](#)].
- [26] A. Sen, *Black hole entropy function, attractors and precision counting of microstates*, *Gen. Rel. Grav.* **40** (2008) 2249 [[arXiv:0708.1270](#)] [[INSPIRE](#)].
- [27] K. Skenderis and P.K. Townsend, *Gravitational stability and renormalization group flow*, *Phys. Lett. B* **468** (1999) 46 [[hep-th/9909070](#)] [[INSPIRE](#)].
- [28] J. de Boer, E.P. Verlinde and H.L. Verlinde, *On the holographic renormalization group*, *JHEP* **08** (2000) 003 [[hep-th/9912012](#)] [[INSPIRE](#)].
- [29] S. de Haro, S.N. Solodukhin and K. Skenderis, *Holographic reconstruction of space-time and renormalization in the AdS/CFT correspondence*, *Commun. Math. Phys.* **217** (2001) 595 [[hep-th/0002230](#)] [[INSPIRE](#)].
- [30] M. Bianchi, D.Z. Freedman and K. Skenderis, *Holographic renormalization*, *Nucl. Phys. B* **631** (2002) 159 [[hep-th/0112119](#)] [[INSPIRE](#)].
- [31] V. Balasubramanian, J. de Boer, M.M. Sheikh-Jabbari and J. Simon, *What is a chiral 2D CFT? And what does it have to do with extremal black holes?*, *JHEP* **02** (2010) 017 [[arXiv:0906.3272](#)] [[INSPIRE](#)].
- [32] E. D'Hoker and P. Kraus, *Magnetic brane solutions in AdS*, *JHEP* **10** (2009) 088 [[arXiv:0908.3875](#)] [[INSPIRE](#)].
- [33] A. Almuhairi, *AdS<sub>3</sub> and AdS<sub>2</sub> magnetic brane solutions*, [arXiv:1011.1266](#) [[INSPIRE](#)].
- [34] A. Almuhairi and J. Polchinski, *Magnetic AdS × R<sup>2</sup>: supersymmetry and stability*, [arXiv:1108.1213](#) [[INSPIRE](#)].
- [35] A. Almheiri, *Magnetic AdS<sub>2</sub> × R<sup>2</sup> at weak and strong coupling*, [arXiv:1112.4820](#) [[INSPIRE](#)].
- [36] S. Jain, N. Kundu, K. Sen, A. Sinha and S.P. Trivedi, *A strongly coupled anisotropic fluid from dilaton driven holography*, [arXiv:1406.4874](#) [[INSPIRE](#)].
- [37] S. Barisch-Dick, G. Lopes Cardoso, M. Haack and S. Nampuri, *Extremal black brane solutions in five-dimensional gauged supergravity*, *JHEP* **02** (2013) 103 [[arXiv:1211.0832](#)] [[INSPIRE](#)].
- [38] G. Dall'Agata and A. Gnecci, *Flow equations and attractors for black holes in N = 2 U(1) gauged supergravity*, *JHEP* **03** (2011) 037 [[arXiv:1012.3756](#)] [[INSPIRE](#)].
- [39] S. Barisch, G. Lopes Cardoso, M. Haack, S. Nampuri and N.A. Obers, *Nernst branes in gauged supergravity*, *JHEP* **11** (2011) 090 [[arXiv:1108.0296](#)] [[INSPIRE](#)].
- [40] S. Barisch-Dick, G.L. Cardoso, M. Haack and Á. Véliz-Osorio, *Quantum corrections to extremal black brane solutions*, *JHEP* **02** (2014) 105 [[arXiv:1311.3136](#)] [[INSPIRE](#)].
- [41] K. Hristov and A. Rota, *Attractors, black objects and holographic RG flows in 5D maximal gauged supergravities*, *JHEP* **03** (2014) 057 [[arXiv:1312.3275](#)] [[INSPIRE](#)].
- [42] I.R. Klebanov, S.S. Pufu and B.R. Safdi, *F-theorem without supersymmetry*, *JHEP* **10** (2011) 038 [[arXiv:1105.4598](#)] [[INSPIRE](#)].
- [43] D.L. Jafferis, I.R. Klebanov, S.S. Pufu and B.R. Safdi, *Towards the F-theorem: N = 2 field theories on the three-sphere*, *JHEP* **06** (2011) 102 [[arXiv:1103.1181](#)] [[INSPIRE](#)].
- [44] A. Sen, *State operator correspondence and entanglement in AdS<sub>2</sub>/CFT<sub>1</sub>*, *Entropy* **13** (2011) 1305 [[arXiv:1101.4254](#)] [[INSPIRE](#)].

- [45] J.D. Brown and M. Henneaux, *Central charges in the canonical realization of asymptotic symmetries: an example from three-dimensional gravity*, *Commun. Math. Phys.* **104** (1986) 207 [[INSPIRE](#)].
- [46] T. Banks and E.J. Martinec, *The renormalization group and string field theory*, *Nucl. Phys. B* **294** (1987) 733 [[INSPIRE](#)].
- [47] J. Hughes, J. Liu and J. Polchinski, *Virasoro-shapiro from Wilson*, *Nucl. Phys. B* **316** (1989) 15 [[INSPIRE](#)].
- [48] V. Balasubramanian, E.G. Gimon, D. Minic and J. Rahmfeld, *Four-dimensional conformal supergravity from AdS space*, *Phys. Rev. D* **63** (2001) 104009 [[hep-th/0007211](#)] [[INSPIRE](#)].
- [49] S. Kachru, X. Liu and M. Mulligan, *Gravity duals of Lifshitz-like fixed points*, *Phys. Rev. D* **78** (2008) 106005 [[arXiv:0808.1725](#)] [[INSPIRE](#)].
- [50] A. Donos and J.P. Gauntlett, *Lifshitz solutions of  $D = 10$  and  $D = 11$  supergravity*, *JHEP* **12** (2010) 002 [[arXiv:1008.2062](#)] [[INSPIRE](#)].
- [51] J. de Boer, J. Manschot, K. Papadodimas and E. Verlinde, *The chiral ring of  $AdS_3/CFT_2$  and the attractor mechanism*, *JHEP* **03** (2009) 030 [[arXiv:0809.0507](#)] [[INSPIRE](#)].
- [52] M. Ammon, M. Gutperle, P. Kraus and E. Perlmutter, *Spacetime geometry in higher spin gravity*, *JHEP* **10** (2011) 053 [[arXiv:1106.4788](#)] [[INSPIRE](#)].
- [53] M.M. Anber and D. Kastor, *The attractor mechanism in Gauss-Bonnet gravity*, *JHEP* **10** (2007) 084 [[arXiv:0707.1464](#)] [[INSPIRE](#)].
- [54] M.M. Anber and D. Kastor, *C-functions in Lovelock gravity*, *JHEP* **05** (2008) 061 [[arXiv:0802.1290](#)] [[INSPIRE](#)].
- [55] M.R. Garousi and A. Ghodsi, *On attractor mechanism and entropy function for non-extremal black holes/branes*, *JHEP* **05** (2007) 043 [[hep-th/0703260](#)] [[INSPIRE](#)].
- [56] A. Castro, J.L. Davis, P. Kraus and F. Larsen, *5D attractors with higher derivatives*, *JHEP* **04** (2007) 091 [[hep-th/0702072](#)] [[INSPIRE](#)].